

Research Article

In vitro shoots multiplication from nodal explants of Sesame (*Sesamum indicum* L.)

Muhammad Asad¹, Nadeem Ahmed¹, Amir Sohail^{2*}, Tanveer Burni¹, Fazal Hadi¹, Rehman Ali¹, Muhammad Nauman Khan³, Abdullah Aziz¹ and Abbas Muhammad¹

1. Department of Botany, University of Peshawar, Khyber Pakhtunkhwa-Pakistan

2. State Key Laboratory of Rice Biology, China National Rice Research Institute, Hangzhou-China

3. Department of Botany, Bacha Khan University Charsadda, KP-Pakistan

*Corresponding author's email: amirsohail@aup.edu.pk

Citation

Muhammad Asad, Nadeem Ahmed, Amir Sohail, Tanveer Burni, Fazal Hadi, Rehman Ali, Muhammad Nauman Khan, Abdullah Aziz and Abbas Muhammad. *In vitro* shoots multiplication from nodal explants of (*Sesamum indicum* L.). Pure and Applied Biology. Vol. 9, Issue 1, pp303-308. <http://dx.doi.org/10.19045/bspab.2020.90035>

Received: 12/01/2019

Revised: 06/09/2019

Accepted: 23/10/2019

Online First: 06/11/2019

Abstract

An effective micro propagation system has been developed efficiently through an enriched culture system for *Sesamum indicum* L. used as a medicinal plant and spicy crop significant from a nutritional point of view. The multiplication of shoot was obtained on Murashige and Skoog medium supplemented with 15 g l⁻¹ sucrose, 8 g l⁻¹ agar and fortified with benzyl amino purine (BAP) in ratios of 0.1 -1 mg / l. The maximum number of young shoots was appeared on MS medium supplemented with 0.5 mg / l benzyl amino purine (BAP). Multiple combinations of harmonics were used for multiple inductions. Developed micro-shoots were shifted to culture medium supplemented with IAA (1.0 mg / l) for efficient rooting. For hardening newly developed shoots were carried out and 71% were survived effectively. This system can be successfully applied for mass propagation of *Sesamum indicum*.

Keywords: BAP; Medium MS; Micro propagation; Multiple shoots; Sesame

Introduction

Sesame (*Sesamum indicum* L.) belongs to the family Pedaliaceae and known as the oldest crop of oilseeds. Sesame cultivated in Afghanistan India, Pakistan, Russia, China, South America and some African countries [1]. Through worldwide sesame cultivated on a total area of over 7.7 million hectares with entire production of 3.3 million tons [2]. Seeds of sesame are vital source of oil and used as spices. It consist of 48-58% of oil, it is very stable due to naturally occurring antioxidants sesamines, sesamol and sesamolol [3]. Sesame oil is used in the

Ayurvedic medicine system. Sesame seeds have antioxidant and antitumor potential. Due to abundant amount of protein the oil cake is used for livestock feed [4].

Sesamum indicum is highly uncooperative to regenerate in vitro conditions. Though, several procedures for micro propagation [5, 6], somatic embryos [7], and indirect regeneration of adventitious [8, 9], were achieved with low frequency. The productivity of *Sesamum indicum* L. is relatively low compared to other oleaginous crops because the cultivation of sesame is limited by poor soils [10, 11]. The present

study shows the results of an experiment to develop an appropriate procedure for the regeneration of multiple shoots in *Sesamum indicum* L.

***In vitro* mass propagation**

Plan tissue culture plays an important role in the study of a growing number of basic and practical programs in plant sciences [12]. Globally tissue culture techniques are used for plant conservation [13]. In this study we propose the conservation and rapid multiplication of Sesame for its use.

Materials and methods

Sesame (*Sesamum indicum* L.) seeds for experimental work have been obtained from the University of Agriculture of Peshawar. Seeds were germinated on Murashige and Skoog (MS) media. Media were pH adjusted before autoclaving at 121°C, 1 atm. The cotton bed method was used with water and MS media for the *in vitro* regeneration of seedlings from seeds. All materials used in cultural work must be free of microbes. This is achieved by one of the following approaches: flame sterilization, 70% ethanol cleaning and other surface sterilants. The tip of shoot and the nodal region of *in vitro* derived seedlings were collected for the induction of multiple shootings. The surface-sterilized seeds were transferred to Murashige and Skoog (MS) medium integrated with 0.8% agar, 3% sucrose and 1% of myoinositol. The media pH was adjusted to 5.7 and the maximum growth of the seedlings was observed and was subsequently sub cultured into the medium containing variable concentrations of BAP, IAA and NAA. The subculture process was carried out with the tip of the shoot and the nodal part of the plants. Therefore, It has been optimized with the maximum shoot lets formed from the sub cultured excised shoot tips. The maximum shooting length and internodes have been determined.

Results and discussion

The seeds that showed *in vitro* regeneration responded successfully. On the 2nd day the seeds were germinated and from the 7th day the maximum growth rate was reached. Plants grown *in vitro* have been used for *in vitro* cultivation studies [3]. The high frequency of germination was recorded on MS basal media similar results were obtained in the *in vitro* regeneration of Sesame by [14].

In *Sesamum indicum* L. both apical and all nodal segments were used as primary explants, the nodal explant is the most commonly used as explant to produce micro shoot [15, 16]. In recent study shoots multiplication from nodal explants was achieved on media supplemented with 0.5 mg / l benzyl amino purine (BAP) and 1.5 mg / l of KIN show high frequency. (84%) Of induction of multiple shoots (Table 3, Fig. 2). 88% Shoots multiplication was achieved by supplement media of 0.5mg/l BAP and 1.5mg/l NAA (Table 2). The same percentage (88%) was also recorded in 0.5mg/l BAP and 1.5mg/l IBA media (Table 4). These results were in agreement with [5, 9, 17].

The well-rooted seedlings were transferred to the potting mix in the ratio of sand: soil: farm manure in a 1: 1 ratio: 1. 70% (Table 1) of the germination percentage was recorded.

In a recent experiment, an effective and reproducible procedure for the multiplication standardization of (*Sesamum indicum* L. CVV SVPR-1) was cultivated and formulated a comparatively efficient and reproducible shoot induction system has been worked out utilizing nodal and apical bud explants. Root induction was performed to increase the efficiency of sesame.

The study revealed that apical shoots and nodal segments for explantation were achieved in *Sesamum indicum* L. On Murashige and Skoog medium integrated with benzylaminopurin (BAP) (Table 5, Fig. 1). (0.5 mg / l) more, Indole acetic acid (IAA)

(1.0 mg / l) *in vitro* cultured micro shoots were successfully developed and transferred

to the potting mix in the sand ratio: soil: courtyard manure (Fig. 3).

Table 1. Effect of BAP on multiplication of shoots per shoot tip cultured on MS media

Hormone BAP(mg/l)	Total No. of explants	No. of explants responded	Percentage of responding cultures	M±SD
0.1	50	24	48%	2.16±1.158
0.2	50	35	70%	2.40±1.130
0.3	50	41	82%	3.31±2.198
0.4	50	38	76%	2.26±1.114
0.5	50	48	96%	4.48±3.292
0.6	50	34	68%	2.98±2.341
0.7	50	29	58%	1.46±1.198
0.8	50	34	68%	2.83±2.106
0.9	50	40	80%	3.09±2.347
1.0	50	32	64%	2.67±2.159




Figure 1. Effect of different concentration of BAP on multiplication of shoots

Table 2. Effect of BAP and NAA on multiplication of shoots per shoot tip as explants

Conc. of BAP (mg/l)	Conc. of NAA (mg/l)	Total No. of explants	Percentage of responding cultures	Percentage of responding cultures	M±SD
0.5	0.5	50	32	64%	3.27±2.742
0.5	1.0	50	38	76%	3.16±2.483
0.5	1.5	50	44	88%	4.45±3.946
0.5	2.0	50	23	46%	2.54±2.06
0.5	2.5	50	18	36%	1.78±1.53




Figure 2. Effect of BAP and KIN on multiplication of shoots per shoot tip as explant

Table 3. Effect of BAP and KIN on multiplication of shoots per shoot tip as explants

BAP (mg/l)	KIN (mg/l)	Total No. of explants	No. of explants responded	Percentage of cultures responding	M±SD
0.5	0.5	50	37	74%	3.48±2.651
0.5	1.0	50	24	48%	3.14±2.63
0.5	1.5	50	42	84%	4.66±3.254
0.5	2.0	50	20	40%	2.87±2.16
0.5	2.5	50	15	30%	2.38±1.59




Figure 3. Effect of IAA on induction of roots in the micro shoots

Table 4. Effect of BAP and IBA on multiplication of shoot per shoot tip as explants

BAP (mg/l)	IBA (mg/l)	Total No. of explants	No. of explants responded	Percentage of responding cultures	M±SD
0.5	0.5	50	25	50%	3.83±2.492
0.5	1.0	50	18	36%	3.36±2.571
0.5	1.5	50	44	88%	4.57±3.635
0.5	2.0	50	27	54%	3.68±2.051
0.5	2.5	50	37	74%	3.24±2.856

Table 5. Effect of IAA on root induction per shoot tip cultured on MS medium

Hormone IAA(mg/l)	Total No. of explants	No. of explants responded	Percentage of responding cultures	M±SD
0.2	50	18	36%	1.65±0.986
0.4	50	23	46%	2.26±2.187
0.6	50	36	72%	4.45±3.685
0.8	50	22	44%	2.98±2.602
1.0	50	45	90%	6.12±5.062
1.2	50	21	42%	3.35±2.517
1.4	50	29	58%	4.43±3.253
1.6	50	18	36%	3.62±2.163
1.8	50	14	28%	2.54±1.967
2.0	50	27	54%	3.62±2.385

Conclusion

Our Present performance determine the culture conditions for micro propagation of sesame plant and concluded that shoot regeneration of *Sesamum indicum* L. were achieved in *in vitro* on MS medium which containing different concentrations of cytokinins and auxins. It is also suggested that high frequency of shoot growth will be achieved by means of BAP, IBA and IAA application. This work will be also provide a helpful tool for genetic transformation and reproductive growth and for others tissue culture studies.

Authors' contributions

Conceived and designed the experiments: N Ahmed, Performed the experiments: M Asad, Analyzed the data: T Burni, F Hadi, R Ali & MN Khan, Contributed materials/ analysis/ tools: A Aziz & A Muhammad, Wrote the paper: A Sohail.

References

1. Michael MND, Joseph PND & Lara P (2005). The Encyclopedia of Healing Foods. Publisher: Atria, 1st Edition.
2. FAOSTAT (2008). <http://faostat.fao.org>.
3. Brar GS & Ahuja KL (1979). Sesame: its culture, genetics, breeding and biochemistry. In: C.P. Malik (ed.). *Annual Review Plant Science*, Kalyani Publishers, New Delhi. 245-313.
4. Osawa T, Kumon H, Namiki M, Kawakishi S & Fukuda Y (1990). Antimutagenic heat-stable antioxidants, in *Mutagens and Carcinogens in the Diet*. Edited by MW Pariza, HU Aeschbacher, JS Felton and S Sato. Wiley-Liss, New York. pp 223-238.
5. Rao KR & Vaidyanath K (1997). Induction of multiple shoots from seedling shoot tips of different varieties of *Sesamum*. *Indian J of Plant Physiol* 2: 257-261.
6. Gangopadhyay G, Poddar R & Gupta S (1998). Micropropagation of sesame

- (*Sesamum indicum* L.) by *in vitro* multiple shoot production from nodal explants. *Phytomorphol* 48: 83-90.
7. Jeyamary R & Jayabalan N (1997). Influence of growth regulators on somatic embryogenesis in sesame. *Plant Cell, Tissue and Organ Cult* 49: 67-70.
 8. Taskin KM & Turgut K (1997). *In vitro* regeneration of sesame (*Sesamum indicum* L.). *Turk J of Bot* 21: 15-18.
 9. Younghee K (2001). Effects of BA, NAA, 2, 4-D and AgNO₃ treatments on the callus induction and shoot regeneration from hypocotyl and cotyledon of sesame (*Sesamum indicum* L.). *J of the Korean Soc for Horticult Sci* 42: 70-74.
 10. George L, Bapat VA & Rao PS (1987). *In vitro* multiplication of sesame (*Sesamum indicum*) through tissue culture. *Annual Bot* 60: 17-21.
 11. Evans DA, Sharp WR & Bravo JE (1984). Cell culture methods for crop improvement. In: *Handbook of plant cell culture: crop species*. Eds. WR Sharp, DA Evans, PVA mmirato and Yamada Y Macmillan Publishing Co, New York. 47-68.
 12. González RG, Quiroz K, Carrasco B & Caligari P (2010). Plant tissue culture: Current status, opportunities and challenges. *Cien Inv Agr* 37(3): 5-30.
 13. Kale VP (2005). *In vitro* studies in capsicum (*Capsicum annuum* L.). *Indian J of Genetics* 65(4): 329-330.
 14. Baskaran P & Jayabalan N (2006). *In vitro* mass propagation and diverse callus orientation on *Sesamum indicum* L. - an important oil plant. *J Agric Tech* 2: 259-269.
 15. Loksha R, Shashidhara N & Janagoudar BS (2007). Callus induction and plant regeneration in Sesame (*Sesamum indicum* L.) through direct seeding. *Plant cell Biotech Mol Bio* 8(2): 85-88.
 16. Ganesh R, Sharma A, Joshim A, Nanawati GC & Simlot MM (1993). Callus cultures from sesame seed explants. *Annual Agric Res* 14: 388-391.
 17. Reddy KRK & Bahadur, B (1989). *In vitro* multiplication of castor. Recent advances in genetics and cytogenetics. *Plant Tissue Culture* 4(3): 479-82.